

Nuclear Quadrupole Effects in Deeply Bound Pionic Atoms

N. Nose-Togawa^{a,b)}, S. Hirenzaki^{a)} and K. Kume^{a)}

a) Department of Physics, Nara Women's University, Nara 630, Japan

b) RCNP, Osaka University, Ibaraki, Osaka 567, Japan

Abstract

We have studied nuclear quadrupole deformation effects in deeply bound pionic atoms theoretically. We have evaluated the level shifts and widths of the hyperfine components using the first order perturbation theory and compared them with the effects of neutron skin. We conclude that the nuclear quadrupole deformation effects for deeply bound $1s$ and $2p$ states are very difficult to observe and that the effects could be observed for $3d$ states. We also conclude that the deformation effects are sensitive to the parameters of the pion-nucleus optical potential.

PACS: 33.25.Fs, 36.10.Gv

Keywords: nuclear quadrupole effects, deeply bound pionic atom

1. Introduction

Deeply bound pionic atoms such as $1s$ and $2p$ states in heavy nuclei were predicted to be quasi-stable [1] and were expected to be observed by proper experimental methods [2, 3]. There have been a number of efforts to find these states both experimentally and theoretically [4-10]. Very recently, deeply bound pionic atoms have been discovered by Yamazaki *et al.* [11] using $(d, {}^3\text{He})$ reactions [7, 12]. In the data we clearly identify a peak structure in the energy spectrum of the ${}^3\text{He}$ which is mainly due to the formation of deeply bound $2p$ atomic state with a $2p_{3/2}^{-1}$ neutron hole state. Formation of the deepest $1s$ atomic state is also studied theoretically [13] and will be tried to find experimentally in near future [14]. We can observe the energy levels and widths of the atomic states by knowing the reaction Q -value of the peak position and the peak width of the ${}^3\text{He}$ spectrum. Hence, we can expect to develop precise spectroscopic studies of deeply bound pionic atoms using the reaction with high energy-resolution. Since deeply bound pionic atoms exist very close to nuclear surface [2, 3], and since they can be populated in all heavy nuclei including β unstable ones using the $(d, {}^3\text{He})$ reactions in principle [6], we can expect to obtain new information on the nuclear sur-

face and pion-nucleus interaction from the spectroscopic studies of the deeply bound pionic states.

Nuclear quadrupole deformation effects on pionic processes have been studied both in bound states [15, 16] and scattering states [17]. In the previous works for the shallow bound states [15, 16], the quadrupole effects were studied in order to extract information on the nuclear deformation using experimental pionic atom spectra. In this paper, we study the quadrupole effects for deeply bound pionic states from different point of view. In order to know the property of the deeply bound pionic atoms well and to consider how we can obtain new information from the pionic atoms, we need to evaluate the nuclear deformation effects. If the effects are large enough, we may be able to use them to observe nuclear deformation precisely, and if the effects are small enough, we can just forget them and extract other information like pion-nucleus interaction and/or neutron skin. Since the deeply bound states have been observed clearly only once, our knowledge is very restricted and we need to study their properties in order to develop the spectroscopic study of the deeply bound pionic states. So far the deformation effects were evaluated only for the shallow pionic atoms such as $3d$ and $4f$ in heavy nuclei and were known to cause level splitting and width change of the atomic states [15, 16]. We can expect to have larger deformation effects for deeper bound states since pion exists closer to nucleus and, thus, feels nuclear surface more sensitively.

In this paper we evaluate the nuclear quadrupole deformation effects on the deeply bound pionic atoms to know how important the effects are in pionic atom spectroscopy. We use the first order perturbation theory to evaluate the level shifts and widths for the individual hyperfine components, which is described in section 2. In section 3 we show the numerical results. We summarize this paper in section 4.

2. Quadrupole effects on pionic atoms

In this section we describe briefly our theoretical formalism which is used to evaluate the quadrupole deformation effects on deeply bound pionic atoms. We apply the first order perturbation theory [15].

The nuclear density is defined by the expectation value of the nuclear ground state with spin J and its z-projection $J_z = J$ as,

$$\rho(\mathbf{r}) = \langle JJ | \hat{\rho}(\mathbf{r}) | JJ \rangle, \quad (1)$$

where $\hat{\rho}(\mathbf{r})$ is defined as,

$$\hat{\rho}(\mathbf{r}) = \sum_{i=1}^A \delta(\mathbf{r} - \mathbf{r}_i), \quad (2)$$

with nuclear mass number A . By multipole expansion, we can rewrite the density as,

$$\rho(\mathbf{r}) = \rho_0(r) + \sum_{\text{even } k \geq 2} \sqrt{\frac{2k+1}{16\pi}} \rho_k(r) Y_k^0(\hat{\mathbf{r}}), \quad (3)$$

where for $k \geq 2$

$$\rho_k(r) = \sqrt{\frac{16\pi}{2k+1}} \langle JJ | \sum_{i=1}^A \frac{\delta(r-r_i)}{r^2} Y_k^0(\hat{\mathbf{r}}_i) | JJ \rangle. \quad (4)$$

Here, the $k = 2$ term corresponds to the nuclear quadrupole density which we will consider in this paper. As a practical parametrization, we use the Woods-Saxon form for the nuclear density distribution in the intrinsic frame:

$$\bar{\rho}(\mathbf{r}) = \frac{\rho_N}{1 + \exp(\frac{r-R(\theta)}{a})} \quad (5)$$

with

$$R(\theta) = R(1 + \beta Y_2^0(\theta)), \quad (6)$$

where ρ_N is the normalization constant, β a quadrupole deformation parameter. The radius parameter is determined as $R = 1.2A^{1/3}$ fm, and the diffuseness parameter is fixed to $a = 0.5$ fm. For the small deformation, we expand the density in powers of β and take first two terms as,

$$\bar{\rho}(\mathbf{r}) \approx \bar{\rho}_0(r) + \sqrt{\frac{5}{16\pi}} \bar{\rho}_2(r) Y_2^0(\theta), \quad (7)$$

where

$$\bar{\rho}_0(r) = \frac{\rho_N}{1 + \exp(\frac{r-R}{a})}, \quad (8)$$

and

$$\bar{\rho}_2(r) = \sqrt{\frac{16\pi}{5}} \beta \rho_N \frac{R}{a} \frac{\exp(\frac{r-R}{a})}{(1 + \exp(\frac{r-R}{a}))^2}. \quad (9)$$

The nuclear densities in the laboratory frame $\rho(r)$ and $\rho_2(r)$, which should be used for our calculations for pionic atoms, are related to those in the intrinsic frame as follows,

$$\rho_0(r) = \bar{\rho}_0(r),$$

and

$$\rho_2(r) = \frac{J(2J-1)}{(J+1)(2J+3)} \bar{\rho}_2(r). \quad (10)$$

The β parameter is related to the nuclear spectroscopic quadrupole moment Q through,

$$Q = \int_0^\infty \rho_2^p(r) r^4 dr, \quad (11)$$

where a superscript ' p ' indicates the proton density distribution. We can make similar approximation to the $\rho^2(r)$ terms which appear in pion-nucleus optical potential as,

$$\rho^2(\mathbf{r}) \approx \rho_0^2(r) + 2\sqrt{\frac{5}{16\pi}} \rho_0(r) \rho_2(r) Y_2^0(\theta). \quad (12)$$

We can calculate pion wave functions by solving the Klein-Gordon equation with the Coulomb potential of a finite nuclear size, and with the pion-nucleus optical potential. We include only spherical nuclear densities for the Klein-Gordon equation and evaluate the quadrupole deformation effect by perturbation. We have applied the standard Ericson-Ericson potential [18] with a parameter set determined by Seki *et al.* [19] and Konijn *et al.* [20]. All the details of the calculation are described in Ref. [3].

Using the β parameter, the potentials can be written as

$$V_{opt} = V_{opt}^{(0)} + \beta V_{opt}^{(2)},$$

and

$$V_C = V_C^{(0)} + \beta V_C^{(2)}. \quad (13)$$

The second terms are due to quadrupole deformation and treated as perturbation. Then, we can express the complex energy shift in the following way:

$$\Delta E = \beta [\langle V_{opt}^{(2)} \rangle + \langle \frac{E - V_C^{(0)}}{\mu} V_C^{(2)} \rangle], \quad (14)$$

where E is the complex eigenvalue of the Klein-Gordon equation and μ the reduced mass of pion. The terms coming from Coulomb and optical potential contribute to the complex energy shift differently as shown in Eq. (14) since these two potential appear in the Klein-Gordon equation in a different way as in Eq. (17).

Because the Klein-Gordon equation includes the complex optical potential which makes the Hamiltonian non-Hermite and makes the eigen energies complex, we should be careful to the normalization condition to get proper orthonormal wave functions. The Klein-Gordon equation can be rewritten as the following two-component equation [21];

$$[-(\sigma_3 + i\sigma_2) \frac{\nabla^2}{2\mu} + \sigma_3 \mu + (\sigma_3 + i\sigma_2) V_{opt}^{(0)} + V_C^{(0)}] \Psi_i = E_i \Psi_i, \quad (15)$$

where σ_i 's are the Pauli spin matrices and

$$\Psi_i = \frac{1}{2} \begin{pmatrix} (1 + (E - V_C^{(0)})/\mu)\phi_i \\ (1 - (E - V_C^{(0)})/\mu)\phi_i \end{pmatrix}. \quad (16)$$

This equation is equivalent to the Klein-Gordon equation,

$$[-\nabla^2 + \mu^2 + 2\mu V_{opt}^{(0)}]\phi_i = (E_i - V_C^{(0)})^2 \phi_i. \quad (17)$$

From Eq. (15), we get orthonormal condition for Ψ as;

$$\int \tilde{\Psi}_i^T \sigma_3 \Psi_j d\mathbf{r} = \delta_{ij}, \quad (18)$$

where the $\tilde{\Psi}$ means to take complex conjugate only for the angular part [21]. This provides proper orthonormal condition for the wave functions as follows;

$$\int \tilde{\phi}_i \frac{E_i + E_j - 2V_C^{(0)}}{2\mu} \phi_j d\mathbf{r} = \delta_{ij} \quad (19)$$

where the $\tilde{\phi}$ means to take complex conjugate only for the angular part of the wave function. We have used the wave function, which satisfies Eqs.(17) and (19), to calculate all expectation values in this paper.

Then, we evaluate the deformation effects on pionic atoms using the first order perturbation theory. To calculate the complex energy shift in Eq. (14), we need following expectation values for the deformation effect due to strong interaction:

$$\langle \hat{\rho}(\mathbf{r}) \rangle|_{k=2} = -\frac{5}{8\pi} C(JlF) \frac{l}{2l+3} \int_0^\infty R_{nl}(r) \rho_2(r) R_{nl}(r) r^2 dr, \quad (20)$$

$$\begin{aligned} \langle \nabla \hat{\rho}(\mathbf{r}) \nabla \rangle|_{k=2} &+ \frac{5}{8\pi} C(JlF) \frac{l}{2l+3} \left\{ \int_0^\infty \frac{dR_{nl}(r)}{dr} \rho_2(r) \frac{dR_{nl}(r)}{dr} r^2 dr \right. \\ &+ [l(l+1) - 3] \int_0^\infty R_{nl}(r) \rho_2(r) R_{nl}(r) dr \left. \right\}, \end{aligned} \quad (21)$$

$$\langle \hat{\rho}^2(\mathbf{r}) \rangle|_{k=2} = -\frac{5}{8\pi} C(JlF) \frac{l}{2l+3} \int_0^\infty R_{nl}(r) (2\rho_0(r) \rho_2(r)) R_{nl}(r) r^2 dr, \quad (22)$$

and

$$\begin{aligned} \langle \nabla \hat{\rho}^2(\mathbf{r}) \nabla \rangle|_{k=2} &+ \frac{5}{8\pi} C(JlF) \frac{l}{2l+3} \left\{ \int_0^\infty \frac{dR_{nl}(r)}{dr} (2\rho_0(r) \rho_2(r)) \frac{dR_{nl}(r)}{dr} r^2 dr \right. \\ &+ [l(l+1) - 3] \int_0^\infty R_{nl}(r) (2\rho_0(r) \rho_2(r)) R_{nl}(r) dr \left. \right\}. \end{aligned} \quad (23)$$

For the deformation effects coming from Coulomb interaction, we need the matrix element:

$$\langle \int \frac{\hat{\rho}^p(\mathbf{r}')}{|\mathbf{r} - \mathbf{r}'|} d^3\mathbf{r}' \rangle_{|k=2} = -\frac{C(JlF)}{2} \frac{l}{2l+3} \int_0^\infty r^2 dr \int_0^\infty r'^2 dr' \frac{r_{\leq}^2}{r_{>}^3} R_{nl}(r) R_{nl}(r) \rho_2^p(r'). \quad (24)$$

In above expressions, $R_{nl}(r)$ is the radial wave function of the bound pion and we have used the coefficient $C(JlF)$ defined as,

$$C(JlF) \equiv \frac{3X(X-1) - 4J(J+1)l(l+1)}{2J(2J-1)l(2l-1)}, \quad (25)$$

with

$$X \equiv J(J+1) + l(l+1) - F(F+1), \quad (26)$$

where l is the angular momentum of pionic states, and $F = l \otimes J$ the total angular momentum of the system. Obviously, the quadrupole deformation has no effect on pionic s states within the first order perturbation.

3. Numerical results

In this section, we evaluate the nuclear quadrupole effects on the deeply bound pionic atoms. We show the results obtained with the potential parameter set [a-1] in Table 1 of ref. [19]. In the numerical calculations, we have used the computer program MATOM [19] to obtain eigen energies and wave functions of pionic $2p$ and $3d$ states. As a typical example of the deformed nucleus, we consider ${}_{71}^{175}\text{Lu}_{104}$ whose ground state is $J^\pi = \frac{7}{2}^+$ and quadrupole moment $Q = +3.49(\text{b})$. We fix the density parameters appeared in the previous section as $R = 6.71 \text{ fm}$ and $a = 0.5 \text{ fm}$ which give $\beta = 0.29$. By solving the Klein-Gordon equation with the spherical density, we obtain the binding energies and widths of the pionic $2p$ and $3d$ states which are B.E.($2p$)=4084, $\Gamma(2p) = 258$, B.E.($3d$)=2133 and $\Gamma(3d) = 29$ in unit of keV.

First, we evaluate the quadrupole deformation effects on pionic $2p$ and $3d$ states. In Fig. 1, we show the level shifts and widths of the pionic $2p$ state (a) and $3d$ state (b) as a function of the β parameter. We can see in the Fig. 1 (a) that the $2p$ level splits into three levels with spacing of order 10keV, which is one order of magnitude smaller than the width of the $2p$ state due to the nuclear absorption of bound pion. Thus the quadrupole hyperfine splitting will not change the level structure significantly for this state. The $3d$ state, as shown in Fig. 2 (b), splits into five levels whose spacing is the same order of magnitude as the width of the $3d$ state. In this case, the level splitting could be seen by proper experiments. For example, we could see the levels as small separated peaks on the top of the large peak consist of all $3d$ state contributions in double differential cross section of the $(d, {}^3\text{He})$ reactions, if we can

achieve high energy-resolution like 10 keV. Even with worse energy resolution, we may be able to see the effects by observing the significantly larger width compared to those calculated with spherical nuclear density.

Next, we evaluate the quadrupole effect for the nucleus with neutron skin. This evaluation is much interesting since deeply bound pionic atoms are known to be sensitive to the neutron skin [22] and since the unstable nuclei are expected to have the neutron skin and the deformation simultaneously. To simulate the existence of the neutron skin, we introduce a parameter Δ as,

$$\Delta = R_n - R_p, \quad (27)$$

and vary the Δ by changing the radius of neutron distribution R_n . We show the calculated energies of deeply bound states as a function of Δ in Fig. 2. As shown in the figure, the deepest $1s$ state is most sensitive to the thickness of the neutron skin. We also show the calculated energy levels of $2p$ and $3d$ atomic state as a function of deformation parameter β , in Fig. 3 for $\Delta = 0.5$ fm and in Fig. 4 for $\Delta = 1.0$ fm. These figures show similar behavior as in Fig. 1 and indicate that the deformation effects on pionic states are qualitatively same for nuclei with thick neutron skin.

It is also interesting to compare the magnitude of the deformation effects with neutron skin effects. For this purpose, we consider two cases $[\beta, \Delta(\text{fm})] = [0.5, 0.0]$ and $[0.0, 1.0]$, and compare the calculated energies in each atomic state. The results are followings:

The deepest $1s$ state is very sensitive to the neutron skin. The energy shift and width change due to neutron skin are $|\Delta E| \sim 300$ keV and $|\Delta \Gamma| \sim 900$ keV, while the energy of the $1s$ state is not affected by the deformation within the first order perturbation theory as mentioned at the end of section 2.

The sensitivities of the $2p$ state to neutron skin are $|\Delta E| \sim 200$ keV, $|\Delta \Gamma| \sim 10$ keV. Those to deformation are $|\Delta E| = 10 \sim 40$ keV, $|\Delta \Gamma| = 10 \sim 70$ keV. The deformation effects depend on the total spin F of the system defined in the last section.

The sensitivities of the $3d$ state to neutron skin are $|\Delta E| \sim 20$ keV, $|\Delta \Gamma| \sim 10$ keV. Those to deformation are $|\Delta E| = 0 \sim 20$ keV, $|\Delta \Gamma| \leq 10$ keV.

Before closing this section, we describe numerical results obtained with another set of potential parameter determined by Konijn *et al.* We used the parameter set 'present fit, $\xi = 1$ ' shown in Table 1 of ref. [20]. We have calculated the binding energies and widths of $2p$ and $3d$ state with the spherical nuclear density and obtained the similar results as those with Seki's potential [19]. At $\Delta = 0$ [fm], we obtained B.E.($2p$) = 4107, $\Gamma(2p)$ = 257 and B.E.($3d$) = 2140, $\Gamma(3d)$ = 31 in unit of keV.

Then we evaluated the quadrupole deformation effects with Konijn's potential and compared them with those by Seki's [a-1] potential. For $3d$ state, Konijn's potential gives 40-60% larger quadrupole shifts with the same sign. As for the width, we obtained much larger deformation effects and its signs are opposite to those with

Seki's potential at $\Delta = 0$ and 0.5 fm, while both potentials give the quadrupole widths with the same sign for $\Delta = 1$ fm. For $2p$ state, Konijn's potential gives opposite sign for both the real and the imaginary parts of the quadrupole energy shifts. Concerning the absolute values, we obtain larger real energy shifts and smaller imaginary energy shifts with Konijn's potential for $\Delta = 0$ and 0.5 fm, while for $\Delta = 1$ fm Seki's potential yields larger real energy shifts and smaller imaginary energy shifts.

We have found that the deformation effects are sensitive to the potential parameters because of the following reasons: As was pointed out by Seki and Masutani [19], the potential parameters are correlated with each other. The isoscalar s-wave term is approximately written as $(b_0 + B_0\rho_e)\rho$ where ρ_e is the effective nuclear density. Then the potential parameters with different b_0 and B_0 but with similar $(b_0 + B_0\rho_e)$ value yield almost the same level shifts and widths for the pionic atom for spherical nuclei. This is the same for p-wave terms. In fact, the real part of the potential parameters by Seki [a-1] and Konijn [$\xi = 1$] are quite different but the numerical values $(b_0 + B_0\rho_e)$ are almost the same. For the deformation effects, however, the different combination of the potential parameter appear as $< (b_0 + 2B_0\rho)\rho_2 > \sim (b_0 + 2B_0\rho_e) < \rho_2 >$ and these are considerably different for the potentials by Seki [a-1] and Konijn. Thus, the potentials which yield the similar results for the level shifts and widths of the spherical nuclei possibly give different quadrupole contribution. Furthermore, there are large cancellation between the deformation effects coming from the strong and Coulomb interaction, so that the moderate changes of the strong quadrupole effects gives the large modification to the total deformation effects. Also, because of the complex pion wave function, the Coulomb term has contributions to real and imaginary energy shifts. These are the reasons why the potentials by Seki and Konijn predicts fairly different quadrupole effects for deeply bound pionic atoms. These situation does not happen for the quadrupole effects in shallow pionic states, because the Coulomb contribution dominates and there are no large cancellation between strong and Coulomb contributions.

4. Summary

We have evaluated the nuclear quadrupole deformation effects on deeply bound pionic atoms. The nuclear deformation induces the hyperfine splitting and affects the widths of the pionic states, which are expected to be larger for deeper states since pion exists closer to the nucleus. We are very much interested in knowing the properties of the deeply bound pionic atoms for deformed nuclei and to find out the best atomic state - nucleus combinations for obtaining new information clearly.

We found that the level shifts and width changes due to the quadrupole deformation are too small to observe for deeply bound $1s$ and $2p$ states. For $2p$ state the natural width of the state due to pion absorption is one order of magnitude larger than the level shift due to nuclear deformation. There are no quadrupole effects on

pionic $1s$ states in the first order perturbation. Thus, it is very unlikely to observe the quadrupole effect in pionic $1s$ and $2p$ states. For atomic $3d$ state, we have a chance to see the level splitting due to the deformation since the natural width of the $3d$ state is only around 30 keV, which is comparable to the level shift due to the quadrupole deformation. Thus, they may be observed with proper experimental technique like $(d, {}^3\text{He})$ reaction [7] with high energy-resolution.

We have also studied the quadrupole effects for nuclei with neutron skin. This is interesting since we expect to have skin and deformation simultaneously in unstable nuclei, where we can produce deeply bound pionic states in principle using the $(d, {}^3\text{He})$ reaction in inverse kinematics [6]. We change the radius of the neutron distribution artificially to simulate the neutron skin. It is found that the existence of the neutron skin does not change the deformation effect significantly. We compare the magnitudes of the both effects and summarize the results in section 3.

We have evaluated the deformation effects using two different optical potential parameters and found that the nuclear quadrupole deformation effects are very sensitive to the potential parameters in deeply bound pionic states. Thus, we can expect to determine the optical potential parameters more precisely if we can observe the nuclear deformation effects on deeply bound pionic atoms systematically.

From the results in this paper, we can conclude that the nuclear quadrupole deformation effects are difficult to observe experimentally for deeply bound pionic states like $1s$ and $2p$ orbits. In order to determine the deformation from the pionic atoms, which could be interesting for unstable nuclei, we need to determine neutron skin thickness (monopole nuclear density) firstly by observing the $1s$ state and then determine the deformation from the observation of $3d$ state with high energy-resolution like 10 keV.

Acknowledgements

One of us (S. H.) acknowledges many discussions and collaborative works on the deeply bound pionic atoms with Prof. H. Toki, Prof. T. Yamazaki, and Prof. R. S. Hayano.

References

- [1] E. Friedman and G. Soff, J. Phys. **G11** (1985) L37.
- [2] H. Toki and T. Yamazaki, Phys. Lett. **B213** (1988) 129.
- [3] H. Toki, S. Hirenzaki, T. Yamazaki and R. S. Hayano, Nucl. Phys. **A501** (1989) 653.
- [4] M. Iwasaki *et al.*, Phys. Rev. **C43** (1991) 1099.
- [5] H. Toki, S. Hirenzaki and T. Yamazaki, Nucl. Phys. **A530** (1991) 679.
- [6] T. Yamazaki, R. Hayano and H. Toki, Nucl. Inst. Methods **A305** (1991) 406.
- [7] S. Hirenzaki, M. Toki and T. Yamazaki, Phys. Rev. **C44** (1991) 2472.
- [8] J. Nieves and E. Oset, Phys. Lett. **B282** (1992) 24.
- [9] T. Yamazaki, Proceedings of Int. Conf. at Wiesbaden, Nucl. Phys. **A553** (1993) 221c.
- [10] N. Matsuoka *et al.*, Phys. Lett. **B359** (1995) 39.
- [11] T. Yamazaki *et al.*, Z. Phys. **A355** (1996) 219.
- [12] H. Toki, S. Hirenzaki and K. Takahashi, Z. Phys. **A356** (1997) 359.
- [13] S. Hirenzaki and H. Toki, Phys. Rev. **C55** (1997) 2719.
- [14] R. S. Hayano and T. Yamazaki, private communication (1996).
- [15] F. Scheck, Nucl. Phys. **B42** (1972) 573.
- [16] B. Olaniyi, A. Shor, S. C. Cheng, G. Dugan and C. S. Wu, Nucl. Phys. **A403** (1982) 572.
- [17] N. Nose, K. Kume and S. Yamaguchi, Phys. Rev. **C50** (1994) 321.
- [18] M. Ericson and T. E. Ericson, Ann. Phys. **36** (1966) 323.
- [19] R. Seki and K. Masutani, Phys. Rev. **C27** (1983) 2799.
- [20] J. Konijn, C.T.A.M.De Laat, A. Taal, and J. H. Koch, Nucl. Phys. **A519** (1990) 773.
- [21] H. Feshbach and F. Villars, Rev. Mod. Phys. **30** (1958) 24.
- [22] H. Toki, S. Hirenzaki and T. Yamazaki, Phys. Lett. **B249** (1990) 391.

Figure Caption

Fig. 1 Binding energies of (a) pionic $2p$ state and (b) $3d$ state as a function of the deformation parameter β defined in the text. Total spin F of each state is indicated in the figures. The width of each state is shown by the error bars.

Fig. 2 Binding energies of pionic $1s$, $2p$ and $3d$ states as a function of neutron skin thickness Δ defined in the text. The width of each state is shown as hatched areas. The hatched area of $3d$ state is obtained by multiplying a factor 10 to the calculated width.

Fig. 3 Same as in Fig. 1 except for neutron thickness parameter $\Delta = 0.5$ fm.

Fig. 4 Same as in Fig. 1 except for neutron thickness parameter $\Delta = 1.0$ fm.

This figure "fig1-1.png" is available in "png" format from:

<http://arXiv.org/ps/nucl-th/9707004v1>

This figure "fig1-2.png" is available in "png" format from:

<http://arXiv.org/ps/nucl-th/9707004v1>

This figure "fig1-3.png" is available in "png" format from:

<http://arXiv.org/ps/nucl-th/9707004v1>

This figure "fig1-4.png" is available in "png" format from:

<http://arXiv.org/ps/nucl-th/9707004v1>